

A key to *Dysoxylum* (Meliaceae) in Australia, with a description of a new species from Far North Queensland

D.J. Mabberley

Abstract

Mabberley, D.J. (Nationaal Herbarium Nederland, University of Leiden, The Netherlands, and National Herbarium of New South Wales, Royal Botanic Gardens Sydney, Mrs Macquaries Road, Sydney 2000, Australia) 2004. A key to *Dysoxylum* (Meliaceae) in Australia, with a description of a new species from Far North Queensland. *Teloepa* 10(3): 725–729. A key to the known 15 species of *Dysoxylum* in mainland Australia is presented. *Dysoxylum pumilum* from Far North Queensland is illustrated and described as new.

Introduction

The continuing delay in publishing my submitted account of Meliaceae for *Flora of Australia* (see Mabberley 1998) has given the opportunity to revise the account of the family in the light of new discoveries. One of these is a new species of *Dysoxylum* Blume from Far North Queensland, here described. Publication of this has been prompted by the imminent appearance of a new edition of William and Wendy Cooper's *Fruits of the Rain Forest* (Cooper & Cooper 1994) in which the species will be figured.

With 15 indigenous species, the genus *Dysoxylum* is the biggest in the family in Australia and, as no modern account of it is accessible, the opportunity is taken to provide a key to the identification of its species here. Descriptions of the 12 species not restricted to Australia were published by the author in Mabberley et al. (1995: 61–133). Descriptions of the remaining two species in mainland Australia, *D. fraserianum* and *D. rufum*, are to be found, in Stanley and Ross (1983: 481) and Taylor and Harden (2002). The indigenous species of Lord Howe Is. (*D. pachyphyllum* Hemsl., endemic) and Norfolk Is. (*D. bijugum* (Lab.) Seem.), not considered here, are described in Green (1994).

Key to the species

Note: The fruits are often characteristic but are rarely well represented in herbaria. Illustrations of most are provided by Cooper and Cooper (1994: tt. 192–203) and are indicated in the key below.

- 1 Leaves decussate
 - 2 Leaflets with scarcely discernible lateral veins; capsule 5–8 cm diam., glabrous
..... *Dysoxylum acutangulum* Miq. subsp. *foveolatum* (Radlk.) Mabb.
northern Australia, Solomon Is. to Sumatra
 - 2* Leaflets with conspicuous lateral veins; capsule c. 2 cm diam., finely hairy
..... *Dysoxylum oppositifolium* F. Muell.
north-east Queensland, New Guinea to Philippines
Cooper and Cooper [1994: t. 198]
- 1* Leaves in spirals

- 3 Leaves in bud stiletto-like
- 4 Leaves 3- or 4-jugate; capsule c. 5 cm long, pyriform, usually strongly veined *in sicco*, lacking rostrum *Dysoxylum latifolium* Benth.
northern Australia, Solomon Is., New Guinea [and Ternate?]
Cooper and Cooper [1994: t. 196]
- 4* Leaves usually 2-jugate; capsule c. 3.5-4 cm long, subspherical, ± veined in sicco, distinctly rostrate *Dysoxylum puuulum* Mabb.
north-east Queensland
- 3* Leaves in bud fist-shaped
- 5 Leaves paripinnate, sometimes with a terminal spike
- 6 Domatia present *Dysoxylum fraserianum* (A. Juss.) Benth.
Queensland, New South Wales
- 6* Domatia absent
- 7 Petals less than 2 mm long
..... *Dysoxylum papuanum* (Merr. & L.M. Perry) Mabb.
north Queensland, New Guinea, Solomon Is.
Cooper and Cooper [1994: t. 199]
- 7* Petals more than 5 mm long
- 8 Bracts and bracteoles 1–1.5 mm long; petals 6–15 mm long; capsule to 2.5 cm diam., foliage not garlic-scented *Dysoxylum klanderi* F. Muell.
north-east Queensland
Cooper and Cooper [1994: t. 195]
- 8* Bracts and bracteoles minute; petals 5–8 mm long; capsule to 7.7 cm diam., foliage garlic-scented *Dysoxylum alliaceum* (Blume) Blume
Queensland, Solomon Is. through Malesia
to peninsular Thailand and Andaman Is.
Cooper and Cooper [1994: t. 192]
- 5* Leaves imparipinnate
- 9 Calyx of free sepals
- 10 Calyx glabrous or subglabrous
..... *Dysoxylum gaudichaudianum* (A. Juss.) Miq.
Queensland to Java, Christmas Is. and Samoa
Cooper and Cooper [1994: t. 194]
- 10* Calyx densely hairy *Dysoxylum rufum* (A. Rich.) Benth.
Queensland, New South Wales
Cooper and Cooper [1994: t. 202]
- 9* Calyx a tube
- 11 Fruit densely hairy *Dysoxylum setosum* (Span.) Miq.
north Queensland, New Guinea, Timor
Cooper and Cooper [1994: t. 203]
- 11* Fruit glabrous
- 12 Flowers on bole and branches, rarely in axils too
- 13 Petals 15–20 mm long *Dysoxylum parasiticum* (Osb.) Kosterm.
Queensland, Solomon Is. through Malesia to Taiwan
Cooper and Cooper [1994: t. 200]
- 13* Petals 8–9 mm long *Dysoxylum pettigrewianum* F.M. Bailey
Queensland, Solomon Is. to Moluccas
Cooper and Cooper [1994: t. 201]
- 12* Flowers in axils

- 14 Flowers 4-merous
 *Dysoxylum mollissimum* Blume subsp. *molle* (Miq.) Mabb.
 Queensland, New South Wales, New Guinea west to Celebes,
 north to Carolines and east to Niue
 Cooper and Cooper [1994: t. 197, as *D. muelleri*]
- 14* Flowers 5-merous *Dysoxylum arborescens* (Blume) Miq.
 Queensland, Vanuatu and Malesia north to Taiwan
 Cooper and Cooper [1994: t. 193]

Dysoxylum pumilum, a new species from Far North Queensland

Dysoxylum pumilum Mabb., species nova

Arbor pumila, gracilis, foliis saepissime bijugatis; *Dysoxylum*, *D. latifolium* Benth. affine, foliolis majoribus forte acuminatis, inflorescentiis longioribus, fructibus subsphericis, rostratis, seminibus majoribus, distincta est.

Typus: Australia, Queensland, Cook, SFR 755, Gosschalk L.A., 17°25'S, 145°47' E, 360 m, B. Hyland 8792, 18 May 1976, fls (holo QRS; iso NSW).

Small understorey tree to 8 m; bole to 8 cm diam. Bark smooth; inner bark white to yellow or brown with no latex. Leafy twigs 3 mm diam., glabrous; apical buds stiletto-like. Leaves 14–25 cm long, 2 (3)-jugate, with terminal spike or its scar, dark glossy green adaxially, alternate (or subopposite); petiole 7.5–10 cm, ± angled, subglabrous, drying blackish; leaflets 12–34 × 5–12 cm, ovate to oblong, chartaceous, glabrous, opposite to alternate, bases weakly asymmetric, acute to obtuse, apices acute to acuminate, the acumen to 8 mm long, costae 8–14 on each side, pale and prominent abaxially in sicco, arcuate, obscure near margin, intercostals subprominent; petiolules 5–8 mm, weakly swollen. Thyrses axillary, sometimes in axils of undeveloped leaves, 8–35 cm long, subspiciform; axis subglabrous to finely sericeous, bearing cymules of 1–3 flowers, rectangular in bud; bracts c. 0.5 mm, triangular, pubescent, caducous; pedicels 3–4 mm long, pubescent, articulated with short pseudopedicels. Calyx c. 2 mm long, 3–4 mm diam., shallowly cupular, margin weakly 4-toothed. Petals 4, 5–6 mm × c. 2.5–3 mm, oblong, cream. Staminal tube narrowed towards apex, subglabrous, cream, margin very shallowly crenulate to ± dentate; anthers 8, c. 1 mm long, included. Disk c. 1 mm tall, glabrous, margin undulate. Ovary finely pubescent, 4-locular; style pubescent in proximal 2/3; stylehead subcapitate. Capsule c. 3.5–4 cm long, 3.5–4 cm diam., subspherical, abruptly rostrate, ± veined in sicco, yellow-green, orange-yellow within. Seeds 2 or 3, c. 2.5–3 cm long, 2–2.5 cm wide, 1–1.5 cm thick, flattened ellipsoid to hemispherical; sarcotesta glossy, orange-brown; cotyledons collateral, dark green.

Other material examined: Queensland, Cook: SFR 755, Gosschalk L.A., EP/34, 17°26'S, 145°46'E, 380 m, *D. Fitzsimon* 222, 30 Nov 1976 (QRS), sterile, and *D. Fitzsimon* 2757, Dec 1976, (QRS), fr.; SFR 755, Gosschalk L.A., 17°24'S, 145°46'E, 380 m, *B. Gray* 786, 9 Nov 1977 (NSW, QRS), fr.; Chuklunga [Chuck Lunga] Creek, 17°25'S, 145°46'E, 360 m, A.K. *Irvine* 1939, 11 July 1979 (QRS), fr.; SFR 755, Bartle Frere, Russell River, 17°23'S, 145°47'E, 380 m, *B. Gray* (*Hyland* no. 25903RFFK), 7 Dec 1995 (QRS), buds; idem, *B. Gray* (*Hyland* no. 25904RFFK) (QRS), sterile; idem, *B. Gray* 6468 (QRS), fr., seedlings; SFR 755, Bartle Frere, Gosschalk L.A., 17°24'S, 145°46' E, 400 m *B. Gray* 6765, 30 May 1996 (QRS), fls; ibid, *B. Gray* 6808. 21 June 1996 (NSW – spirit only, QRS), fls, and 6809 (QRS), fr.; Russell River at junction with Chuck Lunga Creek, 17°26'S, 145°46'E, 160 m, W. *Cooper* 1624, 4 Dec 2001 (QRS), sterile.

Distribution and ecology: an understorey tree known only from the above collections from the Bartle Frere area of north-east Queensland. By comparison, the very closely allied *D. latifolium* is a widespread tree of rain forest and semi-deciduous thicket (Port Moresby region of Papua New Guinea) to 1200 m. It is frequently reported as a common canopy tree in New Guinea and the Solomon Is.




Fig. 1. *Dysoxylum pumilum*. a, habit; b, apical bud; c, flower (dissected); d, flower; e, fruit: a, b, Hyland 8792; c, d, Gray 6808 (spirit material); e, Gray 786. Scale bar: a = 8.5 cm; b = 1.5 cm; c = 0.75 cm; d = 1.2 cm; e = 4 cm.

Etymology: The specific epithet refers to the small size of the treelet.

References

- Cooper, W. and Cooper, W.T. (1994) *Fruits of the rain forest. A guide to fruits in Australian tropical rain forests*. (RD Press: Surry Hills, NSW).
- Green, P.S. (1994) *Dysoxylum*. *Fl. Australia* 49 (Oceanic ls. 1): 247–48.
- Mabberley, D.J., Pannell, C.M. & Sing, A.M. (1995) Meliaceae. *Flora Malesiana* I, 12.
- Mabberley D.J. (1998) Notes on Australian Meliaceae. *Telopea* 8: 45–46.
- Stanley, T.D. & Ross, E.M. (1983) *Flora of south-east Queensland*. Vol. 1 (Department of Primary Industry: Brisbane).
- Taylor, M.J. & Harden, G.J. (2002) Meliaceae. Pp. 327–332 in G.J. Harden (ed.), *Flora of New South Wales*. Revised edition. Vol. 2. (University of New South Wales Press: Sydney).

Manuscript received 4 November 2003

Manuscript accepted 24 November 2003

